

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 04/2022

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Flaveric Tablets 20mg

**Active ingredient:**Benproperine phosphate

**Dosage form:**white tablet, diameter: 6.5 mm, thickness: 3.8 mm

**Print on wrapping:**(Face) Flaveric 20mg, PT F11, (Back) フラベリック 20mg



**Effects of this medicine**

This medicine shows antitussive action on the cough center, lung and bronchi.

It is usually used to improve cough symptoms associated with cold, bronchitis, etc.

**Before using this medicine, be sure to tell your doctor and pharmacist**

- If you have previously experienced any allergic reactions (itching, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

**Dosing schedule (How to take this medicine)**

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 1 tablet (20 mg of benproperine) at a time, 3 times a day. The dosage may be adjusted according to the age or symptoms. Strictly follow the instructions.
- Swallow the tablet without chewing, as you may feel numbness in the mouth if you crunch it.
- If you miss a dose, take the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

**Precautions while taking this medicine**

**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include dry mouth, drowsiness, abdominal pain, rash and abnormal hearing (change in hearing, etc.). If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

**Storage conditions and other information**

- Keep out of reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.