

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 09/2021

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: PIMENOL 100mg

Active ingredient: Pirmenol hydrochloride hydrate

Dosage form: white/white capsule, diameter: 5.9 mm, length: 15.7 mm

Print on wrapping: ピメノールカプセル, 100mg, 696, PIMENOL Cap.



Effects of this medicine

This medicine regulates disturbed pulse by acting on the cardiac muscle.

It is usually used to treat tachyarrhythmia (ventricular).

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itching, rash, etc.) to any medicines.
- If you have: atrioventricular block, sinoatrial block, heart failure, angle-closure glaucoma or urinary retention tendency.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 1 capsule (100 mg of pirmenol) at a time, twice a day. However, the dosage may be adjusted according to the age or symptoms. Strictly follow the instructions.
- If you miss a dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- This medicine may cause syncope, dizziness/light headedness/numbness of limbs, etc. Avoid performing dangerous operations, such as driving a car or operating machinery.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include constipation, gastric distress, difficulty in urination (urinary retention, etc.), headache, insomnia, bitterness in the mouth, nausea, dry mouth and palpitation. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- respiratory distress, dizziness, palpitation [heart failure, ventricular fibrillation, ventricular tachycardia, atrioventricular block, sinus arrest, syncope]
- sweating, chills, palpitation [hypoglycemia]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them. Seek advice of your pharmacy or medical institution about proper disposal of the unused medicines.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.