

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 01/2022


The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:PONTAL FINE GRANULE 98.5%

Active ingredient:Mefenamic acid

Dosage form:faintly yellowish white to pale grayish white fine granules

Print on wrapping:



Effects of this medicine

This medicine suppresses prostaglandin synthesis and thereby eases swelling and pain associated with inflammation and relieves fever.

It is usually used to relieve inflammation/swelling after surgery and injury, to relieve inflammation/pain/fever associated with osteoarthritis, lumbar pain, symptomatic neuralgia and headache, and to relieve fever/pain associated with acute upper respiratory tract inflammation.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itching, rash, etc.) to any medicines.
If you have: peptic ulcer, blood abnormality, liver disorder, renal disorder, cardiac dysfunction, aspirin asthma or its history or hypertension.
- If you are a child with flu-induced fever.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- To relieve inflammation/swelling after surgery and injury, and to relieve inflammation, pain and fever associated with osteoarthritis, lumbar pain, symptomatic neuralgia, headache (when other medicines did not work), sinusitis, menstrual pain, pain after delivery and tooth pain: In general, for adults, take 500 mg of the active ingredient at a time for the first dose, and then take 250 mg every 6 hours. The dosage may be adjusted according to the age or symptoms.
To relieve fever/pain associated with acute upper respiratory tract inflammation (including that accompanied by acute bronchitis): In general, for adults, take 500 mg of the active ingredient at a time, as needed. For children, the standard dose is 6.5 mg/kg of the active ingredient at a time, as needed. The dosage may be adjusted according to the age or symptoms. As a general rule, take this medicine up to twice a day. The maximum daily dose is 1,500 mg for adults.
In any case, avoid taking this medicine on an empty stomach as much as possible. This preparation contains 985 mg of the active ingredient in 1 g. Strictly follow the instructions.
- If you miss a dose, take the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- This medicine may cause sleepiness and dizziness. Pay attention when you operate dangerous machinery, such as driving a car.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include gastrointestinal disorder, nausea, diarrhea/loose stools, rash, thrombocytopenic purpura, hives, redness, itching and vomiting blood. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- cold sweat, breathing difficulty, numbness in the limbs [shock, anaphylaxis]
- symptoms of anemia, headache/sore throat, high fever with chills and shivering [hemolytic anemia, agranulocytosis]
- palpitation, dizziness, bleeding tendency [bone marrow aplasia]

- fever, redness/blister of skin/mucous membrane, pharyngeal pain [toxic epidermal necrosis, oculomuocutaneous syndrome]
- decreased urine output, edema, loss of appetite [acute renal disorder, nephrotic syndrome, interstitial nephritis]
- pain and tenderness in the pit of the stomach, vomiting, diarrhea [peptic ulcer, colitis]
- general malaise, loss of appetite, yellowing of the skin and white of the eyes [fulminant hepatitis, liver dysfunction, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.